

A CLASS OF RECONSTRUCTIBLE GRAPHS

TETSUYA HOSAKA

ABSTRACT. In this paper, we give a class of reconstructible graphs.

1. INTRODUCTION

The purpose of this paper is to give a class of reconstructible graphs.

For a graph G and $v \in V(G)$, let $N(v) = \{u \in V(G) \mid uv \in E(G)\}$ and $N[v] = N(v) \cup \{v\}$.

We prove the following theorem.

Theorem 1.1. *Let G be a graph with $V(G) = \{v_1, \dots, v_k, \dots, v_n\}$ ($1 < k < n$). Suppose that*

- (i) $\bigcup_{i=1}^k N[v_i] = V(G)$,
- (ii) $|d(v_i) - d(v_j)| > 1$ for any $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, n\} \setminus \{i\}$
and
- (iii) $N[v_1] \cap \bigcup_{i=2}^k N[v_i] = \emptyset$.

Then G is reconstructible.

2. PROOF OF THE THEOREM

Let G be a graph with $V(G) = \{v_1, \dots, v_k, \dots, v_n\}$ ($1 < k < n$). Suppose that

- (i) $\bigcup_{i=1}^k N[v_i] = V(G)$,
- (ii) $|d(v_i) - d(v_j)| > 1$ for any $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, n\} \setminus \{i\}$
and
- (iii) $N[v_1] \cap \bigcup_{i=2}^k N[v_i] = \emptyset$.

Let G' be a graph with $V(G') = \{v'_1, \dots, v'_n\}$ such that $G - v_j \cong G' - v'_j$ for any $j \in \{1, \dots, n\}$.

Here we note that $d(v_j) = d(v'_j)$ for any $j \in \{1, \dots, n\}$.

Date: July 11, 2004.

Key words and phrases. the reconstruction conjecture, reconstructible graphs.

Partly supported by the Grant-in-Aid for Scientific Research, The Ministry of Education, Culture, Sports, Science and Technology, Japan, (No. 15740029).

Let $f_j : G - v_j \rightarrow G' - v'_j$ be an isomorphism for each $j \in \{1, \dots, n\}$.

Lemma 2.1. *For each $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, n\} \setminus \{i\}$,*

- (1) $f_j(v_i) = v'_i$ and
- (2) $v_i v_j \in E(G)$ if and only if $v'_i v'_j \in E(G')$.

Proof. (1) We obtain that $f_j(v_i) = v'_i$ from (ii).

(2) Suppose that $v_i v_j \in E(G)$. Then $d(v_i; G - v_j) = d(v_i) - 1$, where $d(v_i; G - v_j)$ is the degree of v_i in $G - v_j$. Here either $d(v'_i; G' - v'_j) = d(v'_i)$ or $d(v'_i; G' - v'_j) = d(v'_i) - 1$. Since $G - v_j \cong G' - v'_j$, $d(v'_i; G' - v'_j) = d(v'_i) - 1$ by (ii). This means that $v'_i v'_j \in E(G')$. Thus if $v_i v_j \in E(G)$ then $v'_i v'_j \in E(G')$. By the same argument, we also can show the reverse. $\square$

Lemma 2.2.

- (i') $\bigcup_{i=1}^k N[v'_i] = V(G')$.
- (ii') $|d(v'_i) - d(v'_j)| > 1$ for any $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, n\} \setminus \{i\}$.
- (iii') $N[v'_1] \cap \bigcup_{i=2}^k N[v'_i] = \emptyset$.

Proof. (ii') is obvious because $d(v_j) = d(v'_j)$ for any $j \in \{1, \dots, n\}$.

(i') Let $j \in \{1, \dots, n\}$. If $j \leq k$ then $v'_j \in N[v'_j]$ and $v'_j \in \bigcup_{i=1}^k N[v'_i]$. Suppose that $k + 1 \leq j$. Since $v_j \in V(G) = \bigcup_{i=1}^k N[v_i]$ by (i), there exists $i_0 \in \{1, \dots, k\}$ such that $v_j \in N[v_{i_0}]$. Then $v_{i_0} v_j \in E(G)$. By Lemma 2.1 (2), $v'_{i_0} v'_j \in E(G')$, i.e., $v'_j \in N[v'_{i_0}] \subset \bigcup_{i=1}^k N[v'_i]$. Thus $\bigcup_{i=1}^k N[v'_i] = V(G')$.

(iii') Suppose that there exists $j \in \{1, \dots, n\}$ such that $v'_j \in N[v'_1] \cap \bigcup_{i=2}^k N[v'_i]$. Then $v'_j \in N[v'_1] \cap N[v'_{i_0}]$ for some $i_0 \in \{2, \dots, k\}$. This means that $v'_1 v'_j \in E(G')$ and $v'_{i_0} v'_j \in E(G')$. By Lemma 2.1 (2), $v_1 v_j \in E(G)$ and $v_{i_0} v_j \in E(G)$. Hence $v_j \in N[v_1] \cap N[v_{i_0}]$. This contradicts (iii). Thus $N[v'_1] \cap \bigcup_{i=2}^k N[v'_i] = \emptyset$. $\square$

Now $f_1 : G - v_1 \rightarrow G' - v'_1$ is an isomorphism. We show that the map $f : G \rightarrow G'$ defined by $f(v_1) = v'_1$ and $f(v_j) = f_1(v_j)$ for each $j \in \{2, \dots, n\}$ is an isomorphism. It is sufficient to prove that for each $j \in \{2, \dots, n\}$, $v_1 v_j \in E(G)$ if and only if $v'_1 f_1(v_j) \in E(G')$. Here we note that $v_1 v_j \in E(G)$ if and only if $v_j \in N(v_1)$, and $v'_1 f_1(v_j) \in E(G')$ if and only if $f_1(v_j) \in N(v'_1)$. Hence we show that $f_1(N(v_1)) = N(v'_1)$. For each $i \in \{2, \dots, k\}$, by (iii) and (iii'),

$$\begin{aligned} f_1(N[v_i]) &= f_1(N[v_i; G - v_1]) = N[f_1(v_i); G' - v'_1] \\ &= N[v'_i; G' - v'_1] = N[v'_i]. \end{aligned}$$

Hence

$$\begin{aligned}
f_1(N(v_1)) &= f_1(V(G - v_1) - \bigcup_{i=2}^k N[v_i]) \\
&= V(G' - v'_1) - \bigcup_{i=2}^k f_1(N[v_i]) \\
&= V(G' - v'_1) - \bigcup_{i=2}^k N[v'_i] \\
&= N(v'_1)
\end{aligned}$$

by (i), (iii), (i') and (iii'). Thus $f_1(N(v_1)) = N(v'_1)$ and $f : G \rightarrow G'$ is an isomorphism. Therefore G is reconstructible.

REFERENCES

- [1] F.S. Mulla, *A class of graphs for which the Ulam conjecture holds*, Discrete Math. **22** (1978), 197–198.

DEPARTMENT OF MATHEMATICS, UTSUNOMIYA UNIVERSITY, UTSUNOMIYA,
321-8505, JAPAN

E-mail address: hosaka@cc.utsunomiya-u.ac.jp